TABLE 1–2 Some Typical Time Intervals			
Time Interval	Seconds (approximate)		
Lifetime of very unstable subatomic particle	10^{-23} s		
Lifetime of radioactive elements	10^{-22} s to 10^{28} s		
Lifetime of muon	10^{-6} s		
Time between human heartbeats	10^0 s (= 1 s)		
One day	10 ⁵ s		
One year	3×10^{7} s		
Human life span	2×10^{9} s		
Length of recorded history	10^{11} s		
Humans on Earth	10^{14} s		
Life on Earth	10 ¹⁷ s		
Age of Universe	10^{18} s		

TABLE 1–3 Some Masses			
Object	Kilograms (approximate)		
Electron	$10^{-30} \mathrm{kg}$		
Proton, neutron	10^{-27}kg		
DNA molecule	$10^{-17} \mathrm{kg}$		
Bacterium	$10^{-15} \mathrm{kg}$		
Mosquito	10^{-5} kg		
Plum	10^{-1} kg		
Human	10^2 kg		
Ship	10^{8} kg		
Earth	$6 \times 10^{24} \text{ kg}$		
Sun	2×10^{30} kg		
Galaxy	10^{41} kg		

Time

The standard unit of time is the second (s). For many years, the second was defined as 1/86,400 of a mean solar day. The standard second is now defined more precisely in terms of the frequency of radiation emitted by cesium atoms when they pass between two particular states. [Specifically, one second is defined as the time required for 9,192,631,770 periods of this radiation.] There are, by definition, 60 s in one minute (min) and 60 minutes in one hour (h). Table 1-2 presents a range of measured time intervals, rounded off to the nearest power of ten.

Mass

The standard unit of mass is the kilogram (kg). The standard mass is a particular platinum-iridium cylinder, kept at the International Bureau of Weights and Measures near Paris, France, whose mass is defined as exactly 1 kg. A range of masses is presented in Table 1-3. [For practical purposes, 1 kg weighs about 2.2 pounds on Earth.]

When dealing with atoms and molecules, we usually use the unified atomic mass unit (u). In terms of the kilogram,

$$1 \text{ u} = 1.6605 \times 10^{-27} \text{ kg}.$$

The definitions of other standard units for other quantities will be given as we encounter them in later Chapters.

Unit Prefixes

In the metric system, the larger and smaller units are defined in multiples of 10 from the standard unit, and this makes calculation particularly easy. Thus 1 kilometer (km) is 1000 m, 1 centimeter is $\frac{1}{100}$ m, 1 millimeter (mm) is $\frac{1}{1000}$ m or $\frac{1}{10}$ cm, and so on. The prefixes "centi-," "kilo-," and others are listed in Table 1-4 and can be applied not only to units of length, but to units of volume, mass, or any other metric unit. For example, a centiliter (cL) is $\frac{1}{100}$ liter (L), and a kilogram (kg) is 1000 grams (g).

Systems of Units

When dealing with the laws and equations of physics it is very important to use a consistent set of units. Several systems of units have been in use over the years. Today the most important is the Système International (French for International System), which is abbreviated SI. In SI units, the standard of length is the meter, the standard for time is the second, and the standard for mass is the kilogram. This system used to be called the MKS (meter-kilogram-second) system.

A second metric system is the cgs system, in which the centimeter, gram, and second are the standard units of length, mass, and time, as abbreviated in the title. The British engineering system takes as its standards the foot for length, the pound for force, and the second for time.

TABLE 1-4 Metric (SI) Prefixes

Prefix	Abbreviation	Value
yotta	Y	10^{24}
zetta	Z	10^{21}
exa	E	10^{18}
peta	P	10^{15}
tera	T	10^{12}
giga	G	10^{9}
mega	M	10^{6}
kilo	k	10^{3}
hecto	h	10^{2}
deka	da	10^{1}
deci	d	10^{-1}
centi	c	10^{-2}
milli	m	10^{-3}
micro†	μ	10^{-6}
nano	n	10^{-9}
pico	p	10^{-12}
femto	f	10^{-15}
atto	a	10^{-18}
zepto	Z	10^{-21}
yocto	У	10^{-24}

 $^{\dagger}\mu$ is the Greek letter "mu."

➡ PROBLEM SOLVING

Always use a consistent set of units

SI units